

EGD PREPARATION

EVENING BEFORE YOUR PROCEDURE:

DO NOT EAT ANYTHING AFTER 10 PM. Follow clear liquid diet instructions.

- Things you can drink: water, black coffee, tea, apple juice, white grape juice, soda, Jell- O, popsicles, broth, and Gatorade or other sport drinks.
- **DO NOT** drink milk or creamer of any kind.
- **DO NOT** drink anything colored red or purple.
- **DO NOT** drink alcohol.

DAY OF PROCEDURE:

- **DO NOT** eat prior to procedure.
- **DO NOT** use chewing tobacco or smoke 6 hours prior to procedure.
- **DO NOT** chew gum, mints, etc. 2 hours prior to procedure time.

You may have clear liquids until 2 hours before your procedure time. DO NOT DRINK ANYTHING 2 HOURS PRIOR TO PROCEDURE.

Important:

You are required to have a driver take you home after the procedure. Arranged transportation will be verified at check in.

Continue taking your prescribed medications, unless you are on a blood thinner and have been told to hold these medications by our office.

Please notify your doctor at least two weeks before the procedure if you take Warfarin (Coumadin), Plavix, Eliquis, Xarelto, Pradaxa, or other blood thinners.

If you are taking one of the following: (Ozempic, Byetta, Byureon, Victoza, Tanzeum, Trulicity, Lixumia, Rybelsus, Wegovy, Fu Laimel, Mounjaro, Exenatide, Liraglutide, Albiglutide, Dulaglutide, Lixisenatide, Beinaglutide, Semaglutide, PEG-loxenatide, Tirzepatide see directions below.

- Daily dose- hold on the day of procedure.
- Weekly dose- hold for 7 days prior to procedure.



Procedure results: communicated via email 2-3 weeks after procedure. Please check your email, including junk, for an email from enki EHR.

Please call our prep line directly for questions related to prep instructions 775-433-0480.

Procedure Location: 5250 Kietzke Lane Reno, NV 89511

*Low residue diet, clear liquid diet, frequently asked questions, procedure tips and more can be found at www.digestivehealthreno.com